

**Instructions.** (100 points) You have 60 minutes. Closed book, closed notes, no calculator. *Show all your work* in order to receive full credit.

- (6<sup>pts</sup>) 1. Consider the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + 3y^2}{3x^2 + y^2}.$$

Either show it does not exist, or give strong evidence for suspecting it does.

- (10<sup>pts</sup>) 2. The following table gives some information about a function  $f(x, y)$ :

$(x, y)$	$f$	$f_x$	$f_y$
$(-1, 3)$	3	2	-1
$(0, 1)$	-5	-1	3
$(3, 4)$	1	4	-2

- (a) (5 pts) Use the chain rule to compute  $\frac{dg}{dt}(0)$  where:

$$g(t) = f(t^2 - t + 3, 2e^{-3t} + 2).$$

- (b) (5 pts) Give an equation for the linear (tangent plane) approximation to  $f$  at the point  $(-1, 3)$ , and use it to estimate  $f(-1.1, 3.2)$ .

(12<sup>pts</sup>) **3.** Evaluate the integral

$$\int_0^4 \int_{\sqrt{y}}^2 e^{(x^3+1)} dx dy$$

fully, by first drawing the region of integration, and then reversing the order of integration.

(12<sup>pts</sup>) **4.** Find and classify (using the Second Derivatives Test) all critical points of


$$f(x, y) = x^2y - 2xy + y^2 - 3y + 1.$$

- (8<sup>pts</sup>) 5. Give an equation for the tangent plane to the surface

$$\frac{xy}{y+z} + e^{-z} \ln(x+2y) = 3$$


at the point  $(3, -1, 0)$ .

- (10<sup>pts</sup>) 6. Use polar coordinates to find the volume of the solid bounded by the cone  $z = \sqrt{x^2 + y^2}$  and the top half of the sphere  $x^2 + y^2 + z^2 = 6$ .



- (16<sup>pts</sup>) 7. A flat triangular plate is bounded by the lines  $y = 2 - 2x$ ,  $y = 2 + 2x$  and the  $x$ -axis, where  $x, y$  are in  $m$ . The mass density is given by

$$\rho(x, y) = y^2 \text{ kg/m}^2.$$




From the symmetry of the plate and the density, you can see that the center of mass of the plate must be on the  $y$ -axis, so  $\bar{x} = 0$ .

- (a) (8 pts) Give an expression involving integrals for  $\bar{y}$ , including appropriate limits of integration.

- (b) (8 pts) The total mass of the plate is  $m = \frac{4}{3}$  kg. Use this to calculate  $\bar{y}$ .

- (10<sup>pts</sup>) 8. Use Lagrange multipliers to find the maximum product of two positive numbers satisfying  $x^2 + y = 6$ .

(16<sup>pts</sup>) **9.** Let  $f(x, y) = x^2y - x + y^2$ .



(a) (5 pts) Compute the directional derivative of  $f$  when moving in the direction of  $-\mathbf{j}$  when you are at the point  $(1, -1)$ . Interpret your result in terms of change in values of  $f$ .

(b) (5 pts) Give the direction and magnitude of maximum decrease of  $f$  when at the point  $(1, -1)$ .

(c) (6 pts) Fully set up bounds and integrand for computing the surface area of  $f$  over the region  $[-1, 2] \times [-2, 1]$ . DO NOT EVALUATE.