

Instructions. You have 120 minutes. Closed book, closed notes, no calculator. **The last page contains some unlabeled theorems from our course.** Show all your work to receive full credit.

1. Consider the points $A = (1, 2, -1)$, $B = (-3, 0, 1)$ and $C = (0, 3, 1)$.

- (a) Give a parameterization of the straight line segment from A to B . Be sure you state what the parameter may range over.

Solution:

$$\begin{aligned}\overrightarrow{AB} &= \langle -3 - 1, 0 - 2, 1 + 1 \rangle = \langle -4, -2, 2 \rangle \Rightarrow \mathbf{r}(t) = \langle 1, 2, -1 \rangle + t \langle -4, -2, 2 \rangle \\ \Rightarrow \quad &\boxed{\mathbf{r}(t) = \langle 1 - 4t, 2 - 2t, 2t - 1 \rangle, \quad 0 \leq t \leq 1}\end{aligned}$$

- (b) Find an equation (*not* a parameterization) for the plane containing A, B, C .

Solution: For the normal vector, we can choose any scalar multiple of $\overrightarrow{AB} \times \overrightarrow{AC}$ for example:


$$\begin{aligned}\overrightarrow{AC} &= \langle 0 - 1, 3 - 2, 1 + 1 \rangle = \langle -1, 1, 2 \rangle \\ \overrightarrow{AB} \times \overrightarrow{AC} &= (-2) \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & -1 \\ -1 & 1 & 2 \end{vmatrix} = -2 \langle 2 + 1, -(4 - 1), 2 + 1 \rangle = -2 \langle 3, -3, 3 \rangle = -6 \langle 1, -1, 1 \rangle \\ \Rightarrow \quad &(x - 1) - (y - 2) + (z + 1) = 0 \quad \text{or} \quad \boxed{x - y + z + 2 = 0}\end{aligned}$$

2. Sketch the region of integration of

$$\int_0^1 \int_{3y}^3 e^{(x^2)} dx dy.$$

Then use your sketch to reverse the order of integration and evaluate the integral.

Solution: The region of integration is:



So switching the order of integration we have:

$$\int_0^1 \int_{3y}^3 e^{(x^2)} dx dy = \int_0^3 \int_0^{\frac{x}{3}} e^{(x^2)} dy dx$$

And we compute:

$$\int_0^3 \int_0^{\frac{x}{3}} e^{(x^2)} dy dx = \int_0^3 \left[y e^{(x^2)} \right]_0^{\frac{x}{3}} dx = \int_0^3 \frac{x}{3} e^{(x^2)} - 0 dx = \left[\frac{e^{(x^2)}}{6} \right]_0^3 = \boxed{\frac{e^9 - 1}{6}}.$$

3. Assume a particle has velocity $\mathbf{v}(t) = (t + 1)\mathbf{i} + 2\sqrt{t}\mathbf{j} + (t - 1)\mathbf{k}$ for $t \geq 1$ with speed measured in m/s.

- (a) Find the time(s) when acceleration is parallel to $2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$.

Solution:

$$\begin{aligned}\mathbf{a}(t) &= \left\langle 1, \frac{1}{\sqrt{t}}, 1 \right\rangle = c \langle 2, 1, 2 \rangle \Rightarrow 1 = 2c \Rightarrow c = \frac{1}{2} \\ \Rightarrow \quad &\frac{1}{\sqrt{t}} = \frac{1}{2} \Rightarrow \sqrt{t} = 2 \Rightarrow \boxed{t = 4 \text{ s}}\end{aligned}$$

- (b) Find the distance traveled from
- $t = 1$
- s to
- $t = 3$
- s.

Solution: Let d be the distance traveled.

$$\begin{aligned}
\|\mathbf{v}(t)\| &= \sqrt{(t+1)^2 + (2\sqrt{t})^2 + (t-1)^2} = \sqrt{t^2 + 2t + 1 + 4t + t^2 - 2t + 1} \\
&= \sqrt{2t^2 + 4t + 2} = \sqrt{2(t+1)^2} = (t+1)\sqrt{2} \\
\Rightarrow d &= \int_1^3 \|\mathbf{v}(t)\| dt = \int_1^3 (t+1)\sqrt{2} dt = \left[\left(\frac{t^2}{2} + t \right) \sqrt{2} \right]_1^3 = \left(\frac{9}{2} + 3 - \frac{1}{2} - 1 \right) \sqrt{2} = \boxed{6\sqrt{2} \text{ m}}
\end{aligned}$$

- (c) Find the position vector
- $\mathbf{r}(t)$
- at all times if
- $\mathbf{r}(1) = 2\mathbf{i} - \frac{1}{2}\mathbf{k}$
- .

Solution:

$$\begin{aligned}
\mathbf{r}(t) - \mathbf{r}(1) &= \int_1^t \mathbf{v}(u) du = \int_1^t \langle u+1, 2\sqrt{u}, u-1 \rangle du = \left[\left\langle \frac{u^2}{2} + u, \frac{4u^{\frac{3}{2}}}{3}, \frac{u^2}{2} - u \right\rangle \right]_1^t \\
\Rightarrow \mathbf{r}(t) - \left\langle 2, 0, -\frac{1}{2} \right\rangle &= \left\langle \frac{t^2}{2} + t, \frac{4t^{\frac{3}{2}}}{3}, \frac{t^2}{2} - t \right\rangle - \left\langle \frac{1}{2} + 1, \frac{4}{3}, \frac{1}{2} - 1 \right\rangle \\
\Rightarrow \mathbf{r}(t) &= \left\langle \frac{t^2}{2} + t, \frac{4t^{\frac{3}{2}}}{3}, \frac{t^2}{2} - t \right\rangle + \left\langle 2 - \frac{3}{2}, 0 - \frac{4}{3}, -\frac{1}{2} + \frac{1}{2} \right\rangle \\
&= \left\langle \frac{t^2}{2} + t, \frac{4t^{\frac{3}{2}}}{3}, \frac{t^2}{2} - t \right\rangle + \left\langle \frac{1}{2}, -\frac{4}{3}, 0 \right\rangle \\
\Rightarrow \mathbf{r}(t) &= \left\langle \frac{t^2}{2} + t + \frac{1}{2}, \frac{4t^{\frac{3}{2}}}{3} - \frac{4}{3}, \frac{t^2}{2} - t \right\rangle
\end{aligned}$$

4. Use Lagrange multipliers to find the extreme values of the function
- $f(x, y) = x^2 - y^2$
- along the parabola
- $x - y^2 = -1$
- .

Solution: Define $g(x, y) = x - y^2 = -1$ for the constraint. Then extreme values will happen when:

$$\nabla f = \lambda \nabla g, \quad g(x, y) = -1 \quad \Rightarrow \quad \langle 2x, -2y \rangle = \lambda \langle 1, -2y \rangle, \quad x - y^2 = -1 \quad \Rightarrow \quad \begin{cases} 2x = \lambda \\ -2y = -2\lambda y \\ x - y^2 = -1 \end{cases}$$

From the second equation, we have two cases:

- if $y = 0$ then from the constraint: $x = -1$ so we have the point $(-1, 0)$;
- if $y \neq 0$ then $\lambda = 1$ and plugging into the first equation we have:

$$2x = 1 \quad \Rightarrow \quad x = \frac{1}{2}$$

which when plugged into the constraint gives you $-y^2 = -\frac{3}{2}$ so $y = \pm\sqrt{\frac{3}{2}}$ and so we have the points $\left(\frac{1}{2}, \pm\sqrt{\frac{3}{2}}\right)$.


We now compute $f(x, y)$ at all the points found above to find the extreme values:

x	y	$f(x, y)$	
-1	0	1	absolute maximum
$\frac{1}{2}$	$\pm\sqrt{\frac{3}{2}}$	$-\frac{5}{4}$	absolute minimum

5. Consider the surface S parametrized by:

$$\mathbf{r}(u, v) = \langle u \cos v, u^2 + 1, u \sin v \rangle \quad \text{for} \quad 0 \leq u \leq 2, \quad 0 \leq v \leq 2\pi$$

in the vector field $\mathbf{F}(x, y, z) = \langle x, yz, y \rangle$ as illustrated below:



- (a) Use Stokes' theorem to compute the circulation of $\mathbf{F}(x, y, z)$ around the oriented boundary curve C of the surface S NOT directly BUT as a surface integral using the given S .

Solution: First we compute:

$$\mathbf{r}_u \times \mathbf{r}_v = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \cos v & 2u & \sin v \\ -u \sin v & 0 & u \cos v \end{vmatrix} = \langle 2u^2 \cos v, -u, 2u^2 \sin v \rangle.$$

Next, we take the curl of the field:

$$\text{curl } \mathbf{F}(x, y, z) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ x & yz & y \end{vmatrix} = \langle 1 - y, 0 - 0, 0 - 0 \rangle = \langle 1 - y, 0, 0 \rangle,$$

and along the surface we have $\text{curl } \mathbf{F}(\mathbf{r}(u, v)) = \langle 1 - (u^2 + 1), 0, 0 \rangle = \langle -u^2, 0, 0 \rangle$. And therefore,

$$\begin{aligned} \iint_S \text{curl } \mathbf{F} \cdot \mathbf{N} \, dS &= \int_0^{2\pi} \int_0^2 \langle -u^2, 0, 0 \rangle \cdot \langle 2u^2 \cos v, -u, 2u^2 \sin v \rangle \, du \, dv = \int_0^{2\pi} \int_0^2 -2u^4 \cos v \, du \, dv \\ &= \left(\int_0^{2\pi} \cos v \, dv \right) \left(\int_0^2 -2u^4 \, du \right) = [\sin v]_0^{2\pi} \left[-\frac{2u^5}{5} \right]_0^2 = \boxed{0} \end{aligned}$$

- (b) Find an equation of the tangent plane to the surface at the point $\left(\frac{1}{2}, 2, \frac{\sqrt{3}}{2}\right)$.

Solution: We first solve for (u, v) such that:

$$\mathbf{r}(u, v) = \left\langle \frac{1}{2}, 2, \frac{\sqrt{3}}{2} \right\rangle \Rightarrow \begin{cases} u \cos v = \frac{1}{2} \\ u^2 + 1 = 2 \\ u \sin v = \frac{\sqrt{3}}{2} \end{cases}.$$

From the second equation we get $u^2 = 1$ so with the restriction in domain of u , that means $u = 1$. Then the first equation becomes $\cos v = \frac{1}{2}$ so $v = \frac{\pi}{3}$ or $v = \frac{5\pi}{3}$, but since $\sin v > 0$ for our point (third equation) that means $v = \frac{\pi}{3}$. Therefore, the normal vector is:

$$\mathbf{r} \left(1, \frac{\pi}{3} \right) = \langle 2u^2 \cos v, -u, 2u^2 \sin v \rangle \Big|_{(1, \frac{\pi}{3})} = \langle 1, -1, \sqrt{3} \rangle.$$

And so the equation of the tangent plane is:


$$\left(x - \frac{1}{2}\right) - (y - 2) + \sqrt{3}\left(z - \frac{\sqrt{3}}{2}\right) = 0 \quad \text{or} \quad \boxed{x - y + z\sqrt{3} = 0}.$$

6. Sketch the two surfaces

$$x^2 + z^2 = 4, \quad y + z = 5$$

and highlight their curve of intersection. Then give a parameterization of that curve.

Solution: We have a cylinder and a slanted plane, so their intersection is elliptic in shape.



7. Find all critical points of the function

$$f(x, y) = x^3 - 6xy + 8y^3$$

and, to the extent possible, determine whether they are local maxima, local minima, or saddle points.

Solution: We have $\nabla f(x, y) = \langle 3x^2 - 6y, -6x + 24y^2 \rangle$ so partials are continuous everywhere and therefore critical points will be where the gradient is null:


$$\begin{aligned} \nabla f(x, y) = \mathbf{0} &\iff \langle 3x^2 - 6y, -6x + 24y^2 \rangle = \langle 0, 0 \rangle \iff \begin{cases} 3x^2 - 6y = 0 \\ -6x + 24y^2 = 0 \end{cases} \\ \Rightarrow \begin{cases} y = \frac{x^2}{2} \\ -x + 4y^2 = 0 \end{cases} &\Rightarrow \begin{cases} y = \frac{x^2}{2} \\ -x + x^4 = 0 \end{cases} \Rightarrow \begin{cases} y = \frac{x^2}{2} \\ x(-1 + x^3) = 0 \end{cases} \end{aligned}$$

From the second equation we get $x = 0$ or $x = 1$ which when plugged into the first equation gives us respectively $y = 0$ or $y = \frac{1}{2}$ so we have two critical points $(0, 0)$ and $\left(1, \frac{1}{2}\right)$. To classify them, we use the Second Derivatives Test:

$$\begin{aligned} f_{xx} &= 6x, \quad f_{yy} = 48y, \quad f_{xy} = -6 \\ \Rightarrow d(x, y) &= f_{xx}f_{yy} - f_{xy}^2 = 288xy - 36 \end{aligned}$$

- $d(0, 0) = 0 - 36 = -36 < 0$, so $\boxed{(0, 0, 0) \text{ is a saddle point ;}}$
- $d(1, 1/2) = 144 - 36 = 108 > 0$, $f_{xx}(1, 1/2) = 6 > 0$ so $\boxed{(1, 1/2) \text{ is a local minimum}}.$

8. Use **cylindrical coordinates** to find the mass of the solid enclosed below by the paraboloid $z = x^2 + y^2 + 1$ and above by the sphere $x^2 + y^2 + z^2 = 5$ if the density function is given by $\rho(x, y, z) = \frac{1}{z^2}$.



Solution: In cylindrical coordinates, the paraboloid is $z = r^2 + 1$, the sphere is $r^2 + z^2 = 5$ and since we have part of the top half, we can solve for $z = \sqrt{5 - r^2}$ and the density function remains $\frac{1}{z^2}$. For bounds in r , note that the shadow of the solid onto the xy -plane is a disk whose radius corresponds to that of the circle of intersection between the paraboloid and the sphere:

$$r^2 + 1 = \sqrt{5 - r^2} \Rightarrow (r^2 + 1)^2 = 5 - r^2 \Rightarrow r^4 + 2r^2 + 1 = 5 - r^2 \Rightarrow r^4 + 3r^2 - 4 = 0 \Rightarrow (r^2 + 4)(r^2 - 1) = 0.$$

So we find $r = 1$ and therefore the mass of the solid is:

$$\begin{aligned} m &= \int_0^{2\pi} \int_0^1 \int_{r^2+1}^{\sqrt{5-r^2}} \frac{1}{z^2} r \, dz \, dr \, d\theta = \left(\int_0^{2\pi} d\theta \right) \left(\int_0^1 \left[-\frac{1}{z} \right]_{r^2+1}^{\sqrt{5-r^2}} r \, dr \right) \\ &= \left[\theta \right]_0^{2\pi} \int_0^1 -\frac{r}{\sqrt{5-r^2}} + \frac{r}{r^2+1} \, dr = 2\pi \left[\sqrt{5-r^2} + \frac{1}{2} \ln(r^2+1) \right]_0^1 \\ &= 2\pi \left(2 + \frac{1}{2} \ln 2 - \sqrt{5} - 0 \right) = \boxed{\pi \left(4 + \ln 2 - 2\sqrt{5} \right)}. \end{aligned}$$

9. Let S be the closed surface that encloses the eighth of the unit ball centered at the origin for which $x \geq 0$, $y \leq 0$ and $z \leq 0$, oriented outward. Use Gauss' Divergence Theorem and **spherical coordinates** to fully SET UP an integral computing the flux out of S of the vector field $\mathbf{F}(x, y, z) = \langle x^2, -2yx, xz \rangle$. DO NOT EVALUATE.

Solution: By the divergence theorem, we can rewrite the flux:

$$\oiint_S \mathbf{F} \cdot d\mathbf{S} = \iiint_Q \operatorname{div} \mathbf{F} \, dV$$

where Q is the solid with boundary S . The divergence of the field is:

$$\operatorname{div} \mathbf{F}(x, y, z) = P_x + Q_y + R_z = 2x - 2x + x = x.$$

Now if we rewrite our solid Q in spherical coordinates, we have $0 \leq \rho \leq 1$ (for the unit ball), $-\frac{\pi}{2} \leq \theta \leq 0$ (for $x \geq 0$, $y \leq 0$), and $\frac{\pi}{2} \leq \phi \leq \pi$ (for $z \leq 0$). And since $x = \rho \cos \theta \sin \phi$ we have:

$$\begin{aligned} \oiint_S \mathbf{F} \cdot \mathbf{N} \, dS &= \iiint_Q \operatorname{div} \mathbf{F} \, dV = \iiint_Q x \, dV = \int_{\frac{\pi}{2}}^{\pi} \int_{-\frac{\pi}{2}}^0 \int_0^1 (\rho \cos \theta \sin \phi) \rho^2 \sin \phi \, d\rho \, d\theta \, d\phi \\ &\Rightarrow \boxed{\oiint_S \mathbf{F} \cdot \mathbf{N} \, dS = \int_{\frac{\pi}{2}}^{\pi} \int_{-\frac{\pi}{2}}^0 \int_0^1 \rho^3 \cos \theta \sin^2 \phi \, d\rho \, d\theta \, d\phi}. \end{aligned}$$

10. Consider the vector field

$$\mathbf{F}(x, y, z) = \langle e^{x-y} - z \sin(xz), z^2 - e^{x-y}, 2yz - x \sin(xz) \rangle.$$

(a) Find a potential function for $\mathbf{F}(x, y, z)$.

Solution: We have that for any potential function f , $\mathbf{F}(x, y, z) = \langle P, Q, R \rangle = \langle f_x, f_y, f_z \rangle$. So,

$$f(x, y, z) = \int P \, dx = \int e^{x-y} - z \sin(xz) \, dx = e^{x-y} + \cos(xz) + C_1(y, z)$$

$$f(x, y, z) = \int Q \, dy = \int z^2 - e^{x-y} \, dy = yz^2 + e^{x-y} + C_2(x, z)$$

$$f(x, y, z) = \int R \, dz = \int 2yz - x \sin(xz) \, dz = yz^2 + \cos(xz) + C_3(x, y)$$

$$\Rightarrow \boxed{f(x, y, z) = e^{x-y} + \cos(xz) + yz^2 (+C)}$$

(b) Use your answer to part (a) to evaluate the work done by \mathbf{F} if a particle follows a helical path from the point $(2, 0, 0)$ to the point $(2, 0, 1)$, spiraling counterclockwise one time around the z -axis.

Solution: By the Fundamental Theorem of Line Integrals,


$$\begin{aligned} W &= \int_C \mathbf{F} \cdot d\mathbf{r} = f(2, 0, 1) - f(2, 0, 0) \\ &= e^{2-0} + \cos(2(1)) + 0 - (e^{2-0} + \cos(2(0)) + 0) \\ &= e^2 + \cos 2 - e^2 - 1 = \boxed{\cos 2 - 1}. \end{aligned}$$

11. Use Green's theorem to compute

$$\int_C (e^{\cos x} - x^2 y) \, dx + (\arctan y + xy) \, dy$$

over the closed curve C made up of the line segment from $(0, 0)$ to $(2, 0)$, then three quarters around the circle $x^2 + y^2 = 4$ until $(0, -2)$ then the line segment back to the origin.

Solution: We verify that C is oriented counterclockwise:



Let $I = \int_C (e^{\cos x} - x^2 y) dx + (\arctan y + xy) dy$. Then by Green's theorem,

$$\begin{aligned} I &= \iint_R (\arctan y + xy)_x - (e^{\cos x} - x^2 y)_y dA = \iint_R y + x^2 dA = \int_0^{\frac{3\pi}{2}} \int_0^2 (r \sin \theta + r^2 \cos^2 \theta) r dr d\theta \\ &= \int_0^{\frac{3\pi}{2}} \int_0^2 r^2 \sin \theta + r^3 \cos^2 \theta dr d\theta = \int_0^{\frac{3\pi}{2}} \left[\frac{r^3}{3} \sin \theta + \frac{r^4}{4} \cos^2 \theta \right]_0^2 d\theta = \int_0^{\frac{3\pi}{2}} \frac{8}{3} \sin \theta + 4 \cos^2 \theta d\theta \\ &= \int_0^{\frac{3\pi}{2}} \frac{8}{3} \sin \theta + 2(1 + \cos 2\theta) d\theta = \left[-\frac{8}{3} \cos \theta + 2\theta + \sin 2\theta \right]_0^{\frac{3\pi}{2}} \\ &= 0 + 3\pi + 0 - \left(-\frac{8}{3} + 0 + 0 \right) = \boxed{3\pi + \frac{8}{3}} \end{aligned}$$

12. Let $f(x, y) = \frac{x}{y^2} + x^2 y$.

- (a) What is the directional derivative of f at $(2, 1)$ when moving towards $(0, 2)$? What does it mean for function values?

Solution:

$$\begin{aligned} \nabla f(x, y) &= \left\langle \frac{1}{y^2} + 2xy, -\frac{2x}{y^3} + x^2 \right\rangle \Rightarrow \nabla f(2, 1) = \langle 1 + 4, -4 + 4 \rangle = \langle 5, 0 \rangle \\ \mathbf{v} &= \langle 0 - 2, 2 - 1 \rangle = \langle -2, 1 \rangle \Rightarrow \mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \left\langle -\frac{2}{\sqrt{5}}, \frac{1}{\sqrt{5}} \right\rangle \\ D_{\mathbf{u}} f(2, 1) &= \nabla f(2, 1) \cdot \mathbf{u} = \langle 5, 0 \rangle \cdot \left\langle -\frac{2}{\sqrt{5}}, \frac{1}{\sqrt{5}} \right\rangle = -\frac{10}{\sqrt{5}} + 0 = \boxed{-2\sqrt{5}} \end{aligned}$$

and so the function values decrease.

- (b) Let $x(s, t) = s^2 t$ and $y(s, t) = 2s - t$. Use the appropriate chain rule to find $\frac{\partial f}{\partial t}$ (no direct substitution). Your final answer should only contain s and t but DO NOT simplify.

Solution:

$$\begin{aligned} \frac{\partial f}{\partial t} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial t} = \left(\frac{1}{y^2} + 2xy \right) (s^2) + \left(-\frac{2x}{y^3} + x^2 \right) (-1) \\ &= \boxed{\frac{s^2}{(2s - t)^2} + 2s^4 t(2s - t) + \frac{2s^2 t}{(2s - t)^3} - s^4 t^2} \end{aligned}$$

SOME FORMULAS FROM THEOREMS IN THE COURSE:

$$\begin{aligned} f(B) - f(A) &= \int_{AB} \nabla f \cdot d\mathbf{r} \\ \oint_{C=\partial R} P dx + Q dy &= \iint_R Q_x - P_y dA \\ \oint_{C=\partial S} \mathbf{F} \cdot d\mathbf{r} &= \iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S} \\ \oint_{S=\partial V} \mathbf{F} \cdot d\mathbf{S} &= \iiint_V \text{div } \mathbf{F} dV \end{aligned}$$